



#5

## SEQUENCE LISTING

<110> INABA, Niro  
HORI, Takeya  
ITO, Satoru

<120> Method for Producing Proteins

<130> 0760-0303P

<140> 10/087,775  
<141> 2002-03-05

<150> JP 2001-60973  
<151> 2001-03-05

<160> 15

<170> PatentIn version 3.1

<210> 1  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Primer CCR3F. Nucleic acid used as forward primer of PCR for cloning human CCR3 gene.

<400> 1

tcgcataatga caacacctact agatacagtt

30

<210> 2  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Primer CCR3R. Nucleic acid used as reverse primer of PCR for cloning human CCR3 gene.

<400> 2

tgcgaattca aacacaatag agagttccgg ctctg

35

<210> 3  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Primer FUT3F. Nucleic acid used as forward primer of PCR for cloning human fucosyltransferase gene.

<400> 3

tgcgaattca tggatccccct gggtgccagcc

30

<210> 4  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Primer FUT3R. Nucleic acid used as reverse primer of PCR for cloning human fucosyltransferase gene.

<400> 4  
tgctcgagt caggtgaacc aagccgctat 30

<210> 5  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Primer gp64F. Nucleic acid used as forward primer of PCR for cloning baculovirus gp64 gene.

<400> 5  
tcgcatatgg taagcgctat tgtttatat 30

<210> 6  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Primer gp64R. Nucleic acid used as reverse primer of PCR for cloning baculovirus gp64 gene.

<400> 6  
tgcgaattca tattgtctat tacggttct 30

<210> 7  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Primer CCR3FE. Nucleic acid used as forward primer of PCR for cloning human CCR3 gene.

<400> 7  
tcggaattca tgacaacctc actagataca 30

<210> 8  
<211> 29  
<212> DNA

<213> Artificial Sequence  
<220>  
<223> Primer CCR3RS. Nucleic acid used as reverse primer of PCR for cloning human CCR3 gene.

<400> 8  
tgcgctcgacc aaacacaata gagagttcc

29

<210> 9  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer GnTVF. Nucleic acid used as forward primer of PCR for cloning human GnTV gene.

<400> 9  
agagtcgaca tggctctt cactccgtgg

30

<210> 10  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer GnTVRXho. Nucleic acid used as reverse primer of PCR for cloning human GnTV gene.

<400> 10  
tgactcgagc tataggcagt ctttgc

26

<210> 11  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer GnTVR17. Nucleic acid used as reverse primer of PCR for cloning human GnTV gene.

<400> 11  
tgaggtaccc tataggcagt ctttgc

26

<210> 12  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer FUT3F1. Nucleic acid used as forward primer of PCR for cloning human NdeI gene.

<400> 12  
tcgcataatgg atccccctggg tgcgagccaag 30

<210> 13  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer FUT3R3. Nucleic acid used as reverse primer of PCR for cloning human XhoI gene.

<400> 13  
atgctcgagt caggtaacc aagccgctat 30

<210> 14  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Intervening peptide region recognized by thrombin.

<400> 14

Leu Val Gly Arg Pro Ser  
1 5

<210> 15  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Intervening peptide region recognized by Factor Xa.

<400> 15

Ile Glu Gly Arg  
1